

**Claim Amendments**

This listing of claims will replace all prior versions, and listings, of claims in the application.

**Listing of Claims**

Claim 1. (Currently Amended) ~~The use~~ A method of cosmetically treating the hair, skin or nails, comprising:

applying to the hair, skin or nails an aqueous polymer dispersion of a cross-linked cationic polymer ~~preparable prepared by polymerization of polymerizing:~~

a) 1 to 99.9 % by weight, based on the total amount of monomers from which the polymer is prepared, of at least one cationic or cationogenic vinyl group-containing monomer selected from the group consisting of N-vinylimidazoles, diallylamines, dialkylaminoalkyl(meth)acrylamides and dialkylaminoalkyl(meth)acrylamides and dialkylaminoalkyl (meth)acrylates,

b) 0 to 99 % by weight, based on the total amount of monomers from which the polymer is prepared, of at least one neutral or basic water-soluble monomer different from monomer (a),

c) 0 to 50 % by weight, based on the total amount of monomers from which the polymer is prepared, of at least one unsaturated acid or one unsaturated anhydride,

d) 0 to 50 % by weight of at least one free-radically copolymerizable monomer different from (a), (b) or (c); and

e) 0.05 to 10 % by weight, based on the total amount of monomers from which the polymer is prepared, of at least one crosslinking monomer having at least two ethylenically unsaturated, nonconjugated double bonds,


wherein the amounts a) to e) are selected such that the resulting polymer, optionally after quaternization or protonation, has an overall positive charge in water in the presence of:

f) 1 to 100 % by weight of the saturation amount in the reaction medium of one or more organic or inorganic salts, whereby 100 % by weight saturation is indicative of that amount of salt which just dissolves in the aqueous medium of the monomer ingredients at the reaction temperature without precipitation, and

g) 0.1 to 30 % by weight, based on the total weight of the dispersion, of at least one water-soluble ~~protective~~ colloid with a composition different from a) to e), and ~~subsequent~~ subsequently at least partial ~~quaternization~~ quaternizing monomer (a) for cases where the monomer (a) is not quaternized, ~~in cosmetics~~.

Claim 2. (Currently Amended) The use method as claimed in claim 1 ~~in~~ , which treats the hair cosmetics.

Claim 3. (Currently Amended) The use method as claimed in claim 1, wherein the free-radically polymerizable vinyl group-containing cationic monomer (a) ~~used~~ is at least one N-vinylimidazole derivative of the formula (I),




~~wherein in which~~ the radicals R<sup>1</sup> to R<sup>3</sup>, independently of one another, are hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or phenyl.

Claim 4. (Withdrawn-Amended) The ~~use~~ method as claimed in claim 1, wherein the free-radically polymerizable vinyl group-containing cationic monomer (a) ~~used~~ is at least one diallylamine derivative of the formula (II),



in which the radical R<sup>4</sup> is C<sub>1</sub>-C<sub>24</sub>-alkyl.

Claim 5. (Withdrawn-Amended) The ~~use~~ method as claimed in claim 1, wherein the free-radically polymerizable vinyl group-containing cationic monomer (a) ~~used~~ is at least one dialkylaminoalkyl(meth)acrylamide and dialkylaminoalkyl (meth)acrylate of the formula (III),



in which  $R^5$  and  $R^6$ , independently of one another, are hydrogen or methyl, Z is a nitrogen atom where  $x=1$  or an oxygen atom where  $x=0$ ,  $R^7$  is a linear or branched  $C_1$ - $C_{24}$ -alkylene radical, and  $R^8$  and  $R^9$ , independently of one another, are a  $C_1$ - $C_{24}$ -alkylene radical.

Claim 6. (Currently Amended) The use method as claimed in claim 1, ~~where~~ wherein the monomer (b) ~~used~~ is at least one N-vinylactam.

Claim 7. (Canceled)

Claim 8. (New) The method as claimed in claim 1, wherein the amount of monomer (b) in the monomer mixture ranges from 0.1 to 10 % by weight.

Claim 9. (New) The method as claimed in claim 8, wherein the amount of monomer (b) in the monomer mixture ranges from 10 % by weight to 95 % by weight.

Claim 10. (New) The method as claimed in claim 1, wherein the water soluble monomer (b) is an N-vinylactam, N-vinylacetamide, N-methyl-N-vinylacetamide, N-vinylformamide, acrylamide, methacrylamide, N,N-dimethylacrylamide, N-methylolmethacrylamide, N-vinylloxazolidone, N-vinyltriazole or an hydroxyalkyl (meth)acrylate.

Claim 11. (New) The method as claimed in claim 1, wherein the crosslinking monomer (e) is an acrylic ester, a methacrylic ester, an allyl ether or a vinyl ether of at least a dihydric alcohol.

Claim 12. (New) The method as claimed in claim 1, wherein the amount of crosslinking monomer (e) in the monomer mixture ranges from 0.07 to 5 % by weight.

Claim 13. (New) The method as claimed in claim 1, wherein the amount of colloid (g) in the monomer mixture ranges from 0.5 to 20 % by weight.

Claim 14. (New) The method as claimed in claim 1, wherein the composition is applied to the hair as a conditioning agent or a thickener.